

Química Cuántica de Sólidos

Tarea 02: Método de Hartree-Fock

Dr. Omar De la Peña Seaman

2 septiembre 2021

Nombre del Estudiante: _____

Problema 1 *Koopmans' theorem*

Demonstrate the Koopman's theorem,

$$E_{HF}(N-1) - E_{HF}(N) \approx -\epsilon_n,$$
$$E_{HF}(N+1) - E_{HF}(N) \approx \epsilon_m,$$

where we have assumed that an electron of the n th orbital has been removed, and that an extra electron has been added to the m th orbital. $E_{HF}(M)$ is the total electronic energy for the M -electron system.

.....

Problema 2 *Hartree-Fock-Roothaan method*

Consider a system with four electrons. These four electrons occupy pairwise two orbitals but have different spin dependence, thus the Slater determinant is,

$$\Phi(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4) = \frac{1}{\sqrt{4!}} \begin{bmatrix} \phi_1(\mathbf{r}_1)\alpha(1) & \phi_1(\mathbf{r}_1)\beta(1) & \phi_2(\mathbf{r}_1)\alpha(1) & \phi_2(\mathbf{r}_1)\beta(1) \\ \phi_1(\mathbf{r}_2)\alpha(2) & \phi_1(\mathbf{r}_2)\beta(2) & \phi_2(\mathbf{r}_2)\alpha(2) & \phi_2(\mathbf{r}_2)\beta(2) \\ \phi_1(\mathbf{r}_3)\alpha(3) & \phi_1(\mathbf{r}_3)\beta(3) & \phi_2(\mathbf{r}_3)\alpha(3) & \phi_2(\mathbf{r}_3)\beta(3) \\ \phi_1(\mathbf{r}_4)\alpha(4) & \phi_1(\mathbf{r}_4)\beta(4) & \phi_2(\mathbf{r}_4)\alpha(4) & \phi_2(\mathbf{r}_4)\beta(4) \end{bmatrix},$$
$$\equiv |\phi_1\alpha, \phi_1\beta, \phi_2\alpha, \phi_2\beta|.$$

Applying the Roothaan approximation and the H-F-R restricted method, the orbitals $\phi_1\alpha$, $\phi_1\beta$, $\phi_2\alpha$, and $\phi_2\beta$ can be expanded in the following four basis functions $\chi_1\alpha$, $\chi_1\beta$, $\chi_2\alpha$, and $\chi_2\beta$, then $N_b = 2$ and:

$$\phi_1(\mathbf{r}) = c_{11}\chi_1(\mathbf{r}) + c_{12}\chi_2(\mathbf{r}),$$
$$\phi_2(\mathbf{r}) = c_{21}\chi_1(\mathbf{r}) + c_{22}\chi_2(\mathbf{r}).$$

where the precise form of χ_p is unimportant right now.

We denote the matrix elements as following,

$$\begin{aligned} \langle \chi_p | \chi_n \rangle &\equiv o_{pn}, \\ \langle \chi_p | \hat{h}_1 | \chi_n \rangle &\equiv h_{1,pn}, \\ \langle \chi_p \chi_q | \hat{h}_2 | \chi_n \chi_r \rangle &\equiv h_{2,pqnr}, \end{aligned}$$

with the corresponding values given on the tables,

p	n	o_{pn}	$h_{1,pn}$	p	q	n	r	$h_{2,pqnr}$
1	1	1.0	-3.0	1	1	1	1	0.5
1	2	-0.5	-2.0	1	1	1	2	0.3
2	2	1.1875	-5.0	1	1	2	2	0.2
				1	2	2	1	0.2
				1	2	1	2	0.4
				1	2	2	2	0.3
				2	2	2	2	0.5

and the matrix elements obey the following,

$$\begin{aligned} o_{pn} &= o_{np}^*, \\ h_{1,pn} &= h_{1,np}^*, \\ h_{2,pqnr} &= h_{2,qprn} = h_{2,nrpn}^* = h_{2,rnqp}^*. \end{aligned}$$

Assuming the following orbitals as an initial guess,

$$\begin{aligned} \phi_1 &= a_1 (0.2\chi_1 + 0.8\chi_2), \\ \phi_2 &= a_2 \left(1.0\chi_1 + \frac{4}{17}\chi_2 \right), \end{aligned}$$

obtain a new full set of orbitals ϕ_1 and ϕ_2 after two iteration steps, and compare the eigenvalues ϵ_l and the coefficients c_{pl} between the different steps and the initial guess.

Hint: Do not forget to apply the normalization condition to the $\{\phi_l\}$ orbital set.

.....