

# Química Cuántica de Sólidos

## Tarea 04: Teoría del funcional de la densidad

Dr. Omar De la Peña Seaman

19 octubre 2021

Nombre del Estudiante: \_\_\_\_\_

### **Problema 1** Thomas-Fermi method

In the Thomas-Fermi model, the kinetic energy functional is approximated to the free-electron gas system, as

$$T[n(\mathbf{r})] = c_1 \int d\mathbf{r} n(\mathbf{r})^{5/3} \quad \forall \quad c_1 = \frac{3}{10} (3\pi^2)^{2/3}.$$

Obtain the value of  $c_1$  under that model.

.....

### **Problema 2** Electron density

If the electronic density can be expressed as,

$$n(\mathbf{r}) = \langle \psi | \hat{n}(\mathbf{r}) | \psi \rangle \quad \forall \quad \hat{n}(\mathbf{r}) = \sum_{i=1}^N \delta(\mathbf{r}_i - \mathbf{r}),$$

demonstrate that  $n(\mathbf{r})$  can be also:

$$n(\mathbf{r}) = N \int d\mathbf{r}_2 d\mathbf{r}_3 \dots d\mathbf{r}_N |\psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N)|^2 \quad \forall \quad \mathbf{x}_\alpha \equiv \mathbf{r}_\alpha, s.$$

Hint:  $\int \delta(\mathbf{r}_1 - \mathbf{r}) f(\mathbf{r}_1) d\mathbf{r}_1 = f(\mathbf{r}).$

.....

### **Problema 3** Spin-scaling method

Considering the Kohn-Sham kinetic-energy funcional of a polarized-system, as:

$$T[n_\uparrow, n_\downarrow] = \sum_i \sum_\sigma \left\langle \psi_i^\sigma \left| -\frac{1}{2} \nabla^2 \right| \psi_i^\sigma \right\rangle,$$

where  $\sigma = \uparrow, \downarrow$  and  $i$  is the particle-label, whith the spin densities given by:

$$n_\sigma(\mathbf{r}) = \sum_i |\psi_i^\sigma(\mathbf{r})|^2,$$

then, demonstrate the following:

1) The kinetic-energy funcional can be expressed as,

$$T[n_\uparrow, n_\downarrow] = \frac{1}{2} T[2n_\uparrow] + \frac{1}{2} T[2n_\downarrow].$$

2) In view that the fractional spin-polarization  $\zeta$  is constant over all space,

$$\zeta = \frac{n_\uparrow - n_\downarrow}{n} \quad \forall \quad n = n_\uparrow + n_\downarrow,$$

then,

$$\begin{aligned} T[n_\uparrow, n_\downarrow] &= \frac{1}{2} \left[ (1 + \zeta)^{5/3} + (1 - \zeta)^{5/3} \right] T_0[n], \\ \forall \quad T_0[n] &= \frac{3}{10} (3\pi^2)^{2/3} \int d\mathbf{r} n^{5/3}. \end{aligned}$$

*Hint:* G.L. Oliver, J.P. Perdew, *Phys. Rev. A* **20**, 397 (1979).

.....

#### Problema 4 Exchange-correlation functional: LDA

For the LDA formulation, demonstrate that the exchange in a polarized system has the following form,

$$\begin{aligned} \epsilon_x(n, \zeta) &= \epsilon_x(n, 0) + [\epsilon_x(n, 1) - \epsilon_x(n, 0)] f_x(\zeta), \\ \forall \quad f_x(\zeta) &= \frac{1}{2} \frac{(1 + \zeta)^{4/3} + (1 - \zeta)^{4/3} - 2}{2^{1/3} - 1}, \end{aligned}$$

where  $n$  and  $\zeta$  are the total density and fractional polarization, respectively.

*Hint:* U. von Barth and L. Hedin, *J. Phys. C: Solid State Phys.* **5**, 1629 (1972).

.....