

Estado Sólido I  
Tarea 3: Modelo del Gas de Electrones Libres

Dr. Omar De la Peña Seaman

18 Febrero 2021

Nombre del Estudiante: \_\_\_\_\_

**Problema 1** *Kinetic energy, pressure, and bulk modulus of an electron gas*

(a) Show that the kinetic energy of a three-dimensional gas of  $N$  free electrons at 0 K is

$$U_0 = \frac{3}{5} N \epsilon_F.$$

(b) Derive a relation connecting the pressure and volume of an electron gas at 0 K.

*Hint:* The result may be written as  $p = (2/3)U_0/V$ .

(c) Show that the bulk modulus  $B = -V(\partial p/\partial V)$  of an electron gas at 0 K is  $B = 5p/3 = 10U_0/9V$ .

.....

**Problema 2** *Chemical potential in two dimensions*

Show that the chemical potential of a Fermi gas in two dimensions is given by:

$$\mu(T) = k_B T \ln [\exp(\pi n \hbar^2 / m k_B T) - 1],$$

for  $n$  electrons per unit area.

*Hint:* The density of orbitals of a free electron gas in two dimensions is independent of energy:  $D(\epsilon) = m/\pi \hbar^2$ , per unit area of specimen.

.....

**Problema 3** *Frequency dependence of the electrical conductivity*

Use the equation  $m(dv/dt + v/\tau) = -eE$  for the electron drift velocity  $v$  and consider an oscillating electric field  $E(t) = Ee^{i\omega t}$ . Then find the following:

(a) The velocity  $v$  as a function of time and frequency.

(b) The conductivity at frequency  $\omega$ , given by:

$$\sigma(\omega) = \sigma(0) \left( \frac{1 + i\omega\tau}{1 + (\omega\tau)^2} \right),$$

where  $\sigma(0) = ne^2\tau/m$ .

.....

**Problema 4** *Cohesive energy of free electron Fermi gas*

We define the dimensionless length  $r_s$  as  $r_0/a_H$ , where  $r_0$  is the radius of a sphere that contains one electron, and  $a_H$  is the Bohr radius  $\hbar^2/e^2m$ .

- (a) Show that the average kinetic energy per electron in a free electron Fermi gas at 0 K is  $2.21/r_s^2$ , where the energy is expressed in rydbergs with  $1 \text{ Ry} = me^4/2\hbar^2$ .
- (b) Show that the coulomb energy of a point positive charge  $e$  interacting with the uniform electron distribution of one electron in the volume of a sphere of radius  $r_0$  (given by  $\rho = -e/V$ ) is  $-3e^2/2r_0$  or  $-3/r_s$  in rydbergs. You can consider the point particle located at the origin of the sphere, and use the following expression for the interaction:

$$V_C^+ = \int \frac{e\rho}{|\mathbf{r}|} d\mathbf{r}.$$

- (c) Show that the coulomb self-energy of the electron distribution of the sphere is  $3e^2/5r_0$  or  $6/5r_s$  in rydbergs, using the following expression:

$$V_C^{self} = \int \frac{q(\mathbf{r})\rho}{|\mathbf{r}|} d\mathbf{r},$$

where  $q(\mathbf{r})$  is the charge of the electron distribution at a given  $\mathbf{r}$ .

- (d) Construct the total energy of the system, by summing the kinetic and both coulomb potential energies. Show that the equilibrium value for  $r_s$  is 2.46, and find the cohesive energy value (the minimum of energy).

.....