

Estado Sólido I
Tarea 1: Estructura Cristalina

Dr. Omar De la Peña Seaman

19 agosto 2021

Nombre del Estudiante: _____

Problema 1 *Properties of basic crystal structures*

- (a) Calculate the first nearest-neighbor distance for the bcc and fcc crystal structures.
- (b) Obtain the packing fraction f_e for the bcc and fcc crystal structures.
- (c) Show that the c/a ratio for an ideal hexagonal close-packed structure is $(8/3)^{1/2} = 1.633$.

.....

Problema 2 *Hexagonal space lattice*

The primitive translation vectors of the hexagonal space lattice may be taken as

$$\mathbf{a}_1 = (3^{1/2}a/2)\hat{\mathbf{x}} + (a/2)\hat{\mathbf{y}}; \quad \mathbf{a}_2 = -(3^{1/2}a/2)\hat{\mathbf{x}} + (a/2)\hat{\mathbf{y}}; \quad \mathbf{a}_3 = c\hat{\mathbf{z}}.$$

- (a) Show that the volume of the primitive cell is $(3^{1/2}/2)a^2c$.
- (b) Show that the primitive translations of the reciprocal lattice are

$$\mathbf{b}_1 = (2\pi/3^{1/2}a)\hat{\mathbf{x}} + (2\pi/a)\hat{\mathbf{y}}; \quad \mathbf{b}_2 = -(2\pi/3^{1/2}a)\hat{\mathbf{x}} + (2\pi/a)\hat{\mathbf{y}}; \quad \mathbf{b}_3 = (2\pi/c)\hat{\mathbf{z}},$$

so that the lattice is its own reciprocal, but with a rotation of axes.

.....

Problema 3 *Interplanar separation and Bragg condition*

Consider a plane hkl in a crystal lattice.

- (a) Prove that the reciprocal lattice vector $\mathbf{G} = h\mathbf{b}_1 + k\mathbf{b}_2 + l\mathbf{b}_3$ is perpendicular to this plane.
- (b) Prove that the distance between two adjacent parallel planes of the lattice is $d(hkl) = 2\pi/|\mathbf{G}|$.

- (c) Show for a simple cubic lattice that $d^2 = a^2/(h^2 + k^2 + l^2)$.
- (d) Show that the diffraction condition $2\mathbf{k} \cdot \mathbf{G} = G^2$ is an analogous statement of the Bragg condition, $2d\sin\theta = n\lambda$.

.....

Problema 4 *Scattering amplitude contributions*

From the scattering amplitude,

$$F = \sum_{\mathbf{G}} \int dV n_{\mathbf{G}} \exp [i(\mathbf{G} - \Delta\mathbf{k}) \cdot \mathbf{r}],$$

show that F is negligibly small when $\Delta\mathbf{k}$ differs significantly from any reciprocal lattice vector \mathbf{G} .

.....

Problema 5 *Structure factor of diamond*

The basis of the diamond crystal structure consists of eight atoms if the cell is taken as the conventional cube.

- (a) Find the structure factor S of this basis.
- (b) Find the zeros of S and show that the allowed reflections of the diamond structure satisfy $v_1 + v_2 + v_3 = 4n$, where all indices are even and n is any integer, or else all indices are odd.

.....