

Física Estadística I  
Tarea 04: Gases Ideales Cuánticos — Gas de Bose-Einstein

Dr. Omar De la Peña Seaman

29 Abril 2020

Nombre del Estudiante: \_\_\_\_\_

**Problema 1**    *Entropy of an ideal gas*

Show that the entropy of an ideal gas in thermal equilibrium is given:

1. In the case of bosons by the following expression,

$$S = k_B \sum_k [\langle \hat{n}_k + 1 \rangle \ln \langle \hat{n}_k + 1 \rangle - \langle \hat{n}_k \rangle \ln \langle \hat{n}_k \rangle].$$

2. In the case of fermions as:

$$S = k_B \sum_k [-\langle 1 - \hat{n}_k \rangle \ln \langle 1 - \hat{n}_k \rangle - \langle \hat{n}_k \rangle \ln \langle \hat{n}_k \rangle].$$

.....

**Problema 2**    *Occupation number's fluctuation*

1. Derive, for all three statistics, the following expressions for the occupation number's fluctuation,  $\sigma_{\hat{n}_k}^2 = \langle \hat{n}_k^2 \rangle - \langle \hat{n}_k \rangle^2$ :

$$\text{Bose-Einstein: } \sigma_{\hat{n}_k}^2|_{BE} = \langle \hat{n}_k \rangle + \langle \hat{n}_k \rangle^2.$$

$$\text{Fermi-Dirac: } \sigma_{\hat{n}_k}^2|_{FD} = \langle \hat{n}_k \rangle - \langle \hat{n}_k \rangle^2.$$

$$\text{Maxwell-Boltzmann: } \sigma_{\hat{n}_k}^2|_{MB} = \langle \hat{n}_k \rangle.$$

2. Show that, quite generally:

$$\sigma_{\hat{n}_k}^2 = \langle \hat{n}_k^2 \rangle - \langle \hat{n}_k \rangle^2 = k_B T \left. \frac{\partial \langle \hat{n}_k \rangle}{\partial \mu} \right|_T.$$

.....

**Problema 3** *Thermodynamic relationships for ideal Bose gas*

1. For the ideal Bose gas it was demonstrated that:

$$\frac{1}{\zeta} \left( \frac{\partial \zeta}{\partial T} \right)_V = - \frac{3}{2T} \frac{g_{3/2}(\zeta)}{g_{1/2}(\zeta)},$$

thus, show the following:

$$\frac{1}{\zeta} \left( \frac{\partial \zeta}{\partial T} \right)_p = - \frac{5}{2T} \frac{g_{5/2}(\zeta)}{g_{3/2}(\zeta)}.$$

2. Hence, also show that,

$$\gamma = \frac{C_p}{C_V} = \frac{(\partial \zeta / \partial T)_p}{(\partial \zeta / \partial T)_V} = \frac{5}{3} \frac{g_{5/2}(\zeta) g_{1/2}(\zeta)}{[g_{3/2}(\zeta)]^2}.$$

3. Check that, as  $T$  approaches  $T_c$  from above, both  $\gamma$  and  $C_p$  diverge as  $(T - T_c)^{-1}$ .

.....

**Problema 4** *Bose-Einstein condensation*

Consider a three-dimensional gas of bosons for which the single-particle energy is given by,

$$\epsilon_{\mathbf{k},n} = \frac{\hbar^2 \mathbf{k}^2}{2m} + \alpha n,$$

where  $\alpha$  is a positive constant and  $n = -j, \dots, j$  is an integer number.

1. Find that the expressions for the pressure  $p$  and the mean number of particles  $N$  in terms of the temperature  $T$  and the fugacity  $\zeta = e^{\beta\mu}$  are:

$$N(T, V, \mu) = \sum_{n=-j}^j \frac{V}{\lambda^3} g_{3/2}(\zeta e^{-\beta\alpha n}) + \frac{\zeta e^{\beta\alpha j}}{1 - \zeta e^{\beta\alpha j}},$$

$$p(T, V, \mu) = \frac{k_B T}{\lambda^3} \sum_{n=-j}^j g_{5/2}(\zeta e^{-\beta\alpha n}).$$

2. Write down the condition for Bose-Einstein condensation, in particular, find that the conditions for  $\zeta$  and  $N_\epsilon^{MAX}$  are given by:

$$\zeta = e^{-\beta\alpha j}, \quad N_\epsilon^{MAX} = \sum_{l=0}^{2j} \frac{V}{\lambda^3} g_{3/2}(e^{-\beta\alpha l}).$$

.....

**Problema 5** *Black-Body thermodynamics*

1. Find expressions for the pressure  $p$ , energy density  $E/V$ , entropy density  $S/V$  and specific heat at constant volume  $C_V$  of black-body radiation in a  $d$ -dimensional cavity at temperature  $T$ :

$$\begin{aligned} p &= a_d I_d (k_B T)^{d+1}, & E/V &= d a_d I_d (k_B T)^{d+1}, \\ S/V &= (d+1) a_d I_d k_B (k_B T)^d, & C_V &= d(d+1) a_d I_d k_B V (k_B T)^d, \end{aligned}$$

where the quantities  $a_d$  and  $I_d$  are given by:

$$a_d = \frac{d-1}{2^{d-1}} \frac{1}{\Gamma(d/2) (\sqrt{\pi} c \hbar)^d}, \quad I_d = \frac{1}{d} \Gamma(d+1) \mathcal{Z}(d+1).$$

2. Evaluate these quantities explicitly for  $d = 3$ .

*Hint:*  $\epsilon_k = c\hbar|\mathbf{k}|$ , where the reciprocal vector  $\mathbf{k}$  is quantized as following:  $\mathbf{k} = (\pi/L)\mathbf{n}$ , and  $\mathbf{n}$  is a  $d$ -dimension vector of positive integers.

.....